

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

Black hole masses in NLS1 galaxies from the X-ray excess variance method

Marek Nikolajuk¹, Paweł Gurynowicz¹, and Bożena Czerny²

Abstract.

We estimate black hole masses in Narrow Line Seyfert 1 (NLS1) galaxies at the basis of their X-ray excess variance. We apply the standard approach appropriate for Broad Line Seyfert 1 (BLS1) galaxies. In general, we find that the obtained masses are by a factor ~ 20 too small to agree with values obtained from other methods (reverberation, stellar dispersion). However, a small subset of our NLS1 objects does not require that multiplication, or the correction factor is less than 4. We find that this subset have a soft X-ray photon index, $\Gamma_{0.1-2.4\text{keV}}$, smaller than 2. We thus postulate that this subclass of NLS1 actually belongs to BLS1.

1. Introduction

NLS1 sources are defined as a subclass of broad line objects at the basis of the optical spectra, as sources with the Full Width at Half Maximum (FWHM) of H β line, produced in Broad Line Region, smaller than 2100 km s^{-1} . In BLS1s FWHM(H β) is $\gtrsim 2100 \text{ km s}^{-1}$. It was later found that NLS1 generally have smaller black hole masses for a given luminosity, and strong soft X-ray excesses (Mathur et al. 2001; Crummy et al. 2006). Soft X-ray photon indices, $\Gamma_{0.1-2.4\text{keV}}$, of NLS1 sources have a mean value 3.1, while BLS1 sources are harder, with a mean of 2.1 (e.g. Boller et al. 1996). It suggests that NLS1 and BLS1 differ with respect to the accretion pattern.

We have considered a sample of 21 NLS1 galaxies (e.g. Mrk 110, 335, 478, Ark 564, NGC 4051, 5506, Ton S180, I Zw 1, PG 1211+143) and a sample of BLS1 galaxies (e.g. NGC 4151, 5548, 7469, 3C 120, F9) studied by Nikolajuk et al. 2006. All sources were monitored both in optical and in X-ray band.

The masses of supermassive black hole for all our galaxies were taken from literature, $M_{\text{BH}}^{\text{lit}}$. They were obtained with the reverberation method (e.g. Peterson et al. 2004) or with the stellar velocity dispersion method (e.g. Woo & Urry 2002). Those masses were used to compare with values obtained from X-ray variability. We have determined the masses using the variance method described by Nikolajuk et al. 2006. It is based on the scaling of the normalized variance of X-ray light curves, σ_{NXS}^2 , measured in range 2-10 keV with the M_{BH} value.

¹Institute of Theoretical Physics, University of Białystok, Lipowa 41, PL-15352 Białystok, Poland. Send requests to M.Nikolajuk: mrk@alpha.uwb.edu.pl

²N. Copernicus Astronomical Center, Polish Academy of Sciences, Bartycka 18, PL-00716 Warsaw, Poland




Figure 1. Left panel: Black hole masses estimated with X-ray variance method versus masses taken from literature. The solid line shows the relation $M_{\text{BH}}^{\text{lit}} = M_{\text{BH}}^{\sigma_{\text{NXS}}^2}$. The dotted line shows the relation $M_{\text{BH}}^{\text{lit}} = 20 M_{\text{BH}}^{\sigma_{\text{NXS}}^2}$. Right panel: The soft X-ray photon index plotted versus the ratio $M_{\text{BH}}^{\text{lit}} / M_{\text{BH}}^{\sigma_{\text{NXS}}^2}$. The dotted lines show how many times $M_{\text{BH}}^{\sigma_{\text{NXS}}^2}$ should be multiplied in order to obtain $M_{\text{BH}}^{\text{lit}}$. The dashed line shows arbitrary border $\Gamma_{0.1-2.4 \text{ keV}} = 2$.

2. Results and a small subset of NLS1 galaxies

Generally, values of $M_{\text{BH}}^{\sigma_{\text{NXS}}^2}$ require a multiplication by some factor. Mean value of this factor is 20 (see e.g. Papadakis, 2004). Surprisingly, this factor for four galaxies (Mrk 42, Mrk 507, Mrk 766, MCG -6-30-15) is smaller than 3 and for HB89 1557+272 is smaller than 4 (Fig. 1). The variability of X-ray light in the 2-10 keV range of those 5 NLS1 galaxies is thus very similar to all BLS1 sources.

We find that this subset have values of soft X-ray photon index, $\Gamma_{0.1-2.4 \text{ keV}}$, less than or equal to ~ 2 . Once again it is behaviour of BLS1 galaxies. Spectra in the soft X-ray range of those 5 NLS1 are similar to spectra of BLS1.

On the other hand when we compare accretion rates, we cannot see difference between those 5 NLS1 and other NLS1 galaxies. This small subset of NLS1 accrete, like other NLS1, with $\dot{m} = \dot{M} / \dot{M}_{\text{Edd}}$ higher than 0.1 or even close to Eddington limit. This is opposite to BLS1 sources, which accrete with $\dot{m} \lesssim 0.1$.

Acknowledgments. This work was partially supported by the grant 1P03D00829 of the Polish State Committee for Scientific Research.

References

- Boller, T., Brandt, W. N., & Fink, H. 1996, A&A, 305, 53
- Crummy, J., Fabian, A. C., Gallo, L., & Ross, R. R., 2006, MNRAS, 365, 1067
- Mathur, S., Kuraszewicz, J., & Czerny, B. 2001, NewA, 6, 321
- Nikołajuk, M., Czerny, B., Ziółkowski, J., & Gierliński, M. 2006, MNRAS, 370, 1534
- Papadakis, I. 2004, MNRAS, 348, 207
- Peterson, B. M., Ferrarese, L., Gilbert, K. M., et al., 2004, ApJ, 613, 682
- Woo, J., & Urry, M. 2002, ApJ, 579, 530